

Some non-finitely presented Lie Algebras

Joseph Abarbanel and Shmuel Rosset

Tel-Aviv University

Ramat-Aviv 69978

Israel

email: rosset@math.tau.ac.il,

yossia@math.tau.ac.il

Abstract

Let L be a free Lie algebra over a field k , I a non-trivial proper ideal of L , $n > 1$ an integer. The multiplier $H_2(L/I^n, k)$ of L/I^n is not finitely generated, and so in particular, L/I^n is not finitely presented, even when L/I is finite dimensional.

1 Introduction

If R is a free associative algebra, over a field, and I is a two sided ideal of R , then Lewin proved [5] that I^2 is not finitely generated (as a 2-sided ideal!) when the algebra R/I is infinite dimensional. In other words, R/I^2 is not finitely presented in this case. On the other hand, it is easy to see that when R is finitely generated and R/I is finite dimensional, so is R/I^2 , and hence I^2 is finitely generated.

Similar behavior is seen in groups. If F is a finitely generated free group, and R is a normal subgroup then R' is *normally* finitely generated if, and only if, F/R is finite. In fact Baumslag, Strebel and Thomson proved [1] a stronger fact. Denoting the m -th member of the lower central series by γ_m , they proved that for $m > 1$ the Schur multiplier of $F/\gamma_m R$, $H_2(F/\gamma_m R, \mathbb{Z})$, is not finitely generated (as an abelian group) if F/R is not finite.

We note that for the three statements

- (a) R is normally finitely generated,

(b) R/R' is finitely generated as a module over $G = F/R$,
(c) $H_2(G, \mathbb{Z})$ is finitely generated as an abelian group
we have (a) $\Rightarrow$ (b) $\Rightarrow$ (c).

In this paper we prove a result of similar nature for Lie algebras.

Theorem. 1.1 *Let L be a free Lie algebra with basis X , over a field k , and I be any non-zero proper ideal of L , then $I' = [I, I]$ is not finitely generated as an ideal. In fact, the “Schur multiplier” of L/I^n , $H_2(L/I^n, k)$, is not finitely generated if $n > 1$, and hence L/I^n is not finitely presented.*

Here I^n denotes I , if $n = 1$, and $[I^{n-1}, I]$ if $n > 1$. Our proof closely follows the lines of [1].

In §2 we define some notations and the Magnus embedding. In §3 we build a mapping from the Schur multiplier into a tensor product of $n - 1$ copies of $U(L/I)$. This is similar to the mapping defined in [1]. In §4 we build a specific isomorphism of Hopf modules, keeping in mind that the enveloping algebra of a Lie algebra is a Hopf algebra. In §5 we employ the mapping and show that the image of the “Schur multiplicator” is not finite dimensional, thus proving the theorem.

We wish to thank Alon Wasserman for his help in §4.

2 Preliminaries and Notations

Let $\mathcal{G}$ be a Lie algebra. We will denote the Lie multiplication of two elements $a, b \in \mathcal{G}$ by $[a, b]$. As we will also be considering the enveloping algebra of $\mathcal{G}$, the multiplication in $U(\mathcal{G})$ will be denoted simply as ab , while the action of an element $l \in U(\mathcal{G})$ on an element $a \in \mathcal{G}$ will be denoted by $a \cdot l$. Note that the action is the adjoint action, so that if $l \in L$ then $a \cdot l = [a, l]$.

Let $\mathcal{G}$ be a Lie algebra over a field k , $U(\mathcal{G})$ its enveloping algebra, $\delta U(\mathcal{G})$ the augmentation ideal of U . Suppose $0 \rightarrow I \rightarrow L \rightarrow \mathcal{G} \rightarrow 0$ is a free presentation of $\mathcal{G}$, where L is the free Lie algebra with basis X . The enveloping algebra, $U(L)$, is therefore a free associative algebra, with basis X , and $\delta U(L)$ is a free $U(L)$ module, with a basis in one-to-one correspondence with X . Note that over a field, if $\mathcal{G} \neq 0$, $U(\mathcal{G})$ is infinite dimensional, and is without zero divisors.

In addition, if $\mathcal{G}$ is a Lie algebra over a field and $U(\mathcal{G})$ is its enveloping algebra, let $U_n(\mathcal{G})$ be the subspace of $U(\mathcal{G})$ spanned by all

the products of at most n factors from $\mathcal{G}$. This gives a well known ascending filtration of $U(\mathcal{G})$, and we can define the *degree* of an element l to be the *least* integer n such that $l \in U_n(\mathcal{G})$. This function has the properties:

- 1) $\deg(a + b) \leq \max\{\deg(a), \deg(b)\}$,
- 2) if $\deg(a) < \deg(b)$ then $\deg(a + b) = \deg(b)$,
- 3) $\deg(ab) = \deg(a) + \deg(b)$.

In particular, if $x \in \mathcal{G}$ is non-zero then the degree of x is 1, so if $x_1, x_2, \dots, x_n \in \mathcal{G}$ are all non-zero then $\deg(x_1 x_2 \cdots x_n) = n$.

Via the adjoint action, I/I' carries the structure of a $U(L)$ module, and I acts trivially. All modules will be right modules. Therefore I/I' is a $U(L/I)$ module in a natural way. There is a well known embedding of $U(L/I)$ modules, the Magnus embedding, described below, of I/I' into $\delta U(L) \otimes_{U(L)} U(L/I)$. This embedding will be denoted by $\phi : I/I' \rightarrow \delta U(L) \otimes_{U(L)} U(L/I)$. The action of L on $\delta U(L) \otimes_{U(L)} U(L/I)$ is by right multiplication in the right hand term.

The embedding can be defined in the following way. First define $\phi : I \rightarrow \delta U(L) \otimes_{U(L)} U(L/I)$ by $\phi(x) = x \otimes 1$. By using the Poincare-Birkhoff-Witt theorem, and the structure it gives to $U(L)$, it can be seen that this is a mapping of $U(L)$ modules, i.e. $\phi(a \cdot l) = \phi(a)l$. First we check the statement for elements of L . If $l \in L$ then $a \cdot l = [a, l]$ and $\phi([a, l]) = [a, l] \otimes 1 = (al - la) \otimes 1 = a \otimes l - l \otimes a$. However, $a = 0$ in $U(L/I)$ so $\phi([a, l]) = a \otimes l = (a \otimes 1)l = \phi(a)l$. Consider now the subalgebra $A = \{u \in U(L) | \phi(x \cdot u) = \phi(x)u \quad \forall x \in I\}$. Since $L \subset A$ then $A = U(L)$, thus ϕ is a $U(L)$ module homomorphism.

It is left to show that $\ker \phi = I'$. If $x \in I'$ then x can be written as $x = \sum [a_i, b_i]$, $a_i, b_i \in I$, so that $\phi(x) = x \otimes 1 = \sum [a_i, b_i] \otimes 1 = \sum (a_i b_i - b_i a_i) \otimes 1 = \sum a_i \otimes b_i - b_i \otimes a_i$. Since $a_i, b_i \in I$ then their images in $U(L/I)$ are 0 so that $\phi(x) = 0$.

Therefore $I' \subset \ker \phi$. On the other hand suppose $x \in \ker \phi$. Since $\delta U(L)$ is a free $U(L)$ module with basis $\{x_i\}$ where x_i is a basis of L as a free Lie algebra, we have $x \otimes 1 = \sum x_i \otimes f_i$, where, since $\phi(x) = 0$, $f_i = 0$ in $U(L/I)$. Let us denote by $\tilde{I}$ the kernel of the mapping $U(L) \rightarrow U(L/I)$, so that $f_i \in \tilde{I}$. But $\tilde{I} = U(L)I = IU(L)$ and thus by the Poincare-Birkhoff-Witt theorem this kernel is a free left and right $U(L)$ module with a basis that is a basis of I as a subalgebra of L . Therefore $f_i = \sum w_{i,j} a_j$ where a_j are a basis of I . It follows that $x = \sum x_i w_{i,j} a_j$. Consider now the image of x , $\bar{x}$, in I/I' . Since I/I' is the commutative Lie algebra with a basis that is a basis of I as a subalgebra of L , then $\bar{x} = \sum \lambda_j a_j$, where $\lambda_j \in k$. In other words

$x = \sum \lambda_j a_j + w, w \in I'$. But since $I' \subset \ker \phi$ then we can assume $x = \sum \lambda_j a_j$. On the other hand $\phi(x) = 0$ so $x = \sum x_i w_{i,j} a_j$. Since $\tilde{I}$ is a free $U(L)$ module with basis a_i we have $\lambda_j = \sum x_i w_{i,j}$, but $x_i \in \delta U(L)$, so $\lambda_j = 0$. Hence $x \in I'$, therefore $\ker \phi = I'$.

Another proof of the fact that $\ker \phi = I'$ can be found in [2] §8, as the Magnus embedding is a special case of the derivations defined there.

Throughout the remainder of this paper I will be a proper non-zero ideal of L , and $n > 1$ will be an integer.

3 An image of $H_2(L/I^n, k)$

Consider $H_2(L/I^n, k)$. It is known (e.g. [7] p.233) that the analogue of the Hopf formula for groups holds for Lie algebras. Therefore

$$H_2(L/I^n, k) = I^n/[I^n, L] = (I^n/I^{n+1}) \otimes_{U(L)} k$$

We know from the Širšov-Witt theorem (see e.g. [6] p.44) that I is a free Lie algebra. Hence I^n/I^{n+1} is, in a natural way, identifiable with the n -th homogeneous component of the free Lie algebra with basis that is a basis of I/I' as a vector space. Since the free Lie algebra of a free module can be embedded in the tensor algebra over this module, the n -th homogeneous component can be embedded into the n -fold tensor product, i.e. I^n/I^{n+1} can be embedded in $\otimes^n I/I'$, where the tensor is over k . Any unadorned tensor product below is to be taken to be over k . We need this embedding to be a $U(L/I)$ module homomorphism, and it is easy to see that this is indeed the case when $U(L/I)$ acts on I^n/I^{n+1} via the adjoint action, and on $\otimes^n I/I'$ diagonally. The module $\otimes^n I/I'$ can again be embedded, through the Magnus embedding, into

$$\otimes^n (\delta U(L) \otimes_{U(L)} U(L/I))$$

Tensoring this with k over L we get a mapping

$$H_2(L/I^n, k) \approx \otimes^n I/I' \otimes_{U(L)} k \rightarrow \otimes^n (\delta U(L) \otimes_{U(L)} U(L/I)) \otimes_{U(L)} k$$

Since $\delta U(L)$ is a free $U(L)$ module, with a basis X that is a basis of L as a Lie algebra, we can define for each $x \in X$ a projection, denoted $p_x : \delta U(L) \otimes_{U(L)} U(L/I) \rightarrow U(L/I)$. We therefore have for

each n -tuple $(x_1, x_2, \dots, x_n) \in X^n$ a mapping $\phi_{x_1, \dots, x_n} := (p_{x_1} \otimes \dots \otimes p_{x_n} \otimes 1) \circ \phi$

$$\phi_{x_1, x_2, \dots, x_n} : H_2(L/I^n, k) \rightarrow \otimes^n U(L/I) \otimes_{U(L)} k$$

Since $I/I' \rightarrow U(L/I) \otimes \delta U(L)$ is an embedding, there exist elements $\alpha \in I/I'$ and $x \in X$ such that under the Magnus embedding and the projection by x the image $a = \phi_x(\alpha)$ is non-zero. These elements will be put to use below.

4 Isomorphism of Hopf modules

As seen in the last section the image of the multiplier lies in $(U(L/I) \otimes U(L/I) \cdots \otimes U(L/I)) \otimes_{U(L)} k$. On the other hand it is well known that the enveloping algebra is a Hopf algebra, and the action with which this module is endowed is consistent with the standard Hopf structure on $U(L/I)$, which is the diagonal action. We shall use the following notation for the structure of Hopf algebras and modules. Let H be a Hopf algebra and M a Hopf module over H . The diagonal mapping of H will be denoted by Δ , and the n -fold application of Δ by Δ_n (by the co-associativity of H the components on which we apply Δ each time do not matter). The co-unit of H will be denoted by ϵ (also sometimes known as the augmentation). The antipode map of H will be denoted by S . The usual action of H on M will be denoted by multiplication on the right, and the co-action of M will be denoted by ρ . If $h \in H$ then $\Delta(h)$ will be written as $\Delta(h) = \sum_{i=1}^l h_1^i \otimes h_2^i$, and $\Delta(h_1^i) = \sum_{j=1}^{l(i)} h_{1,1}^{i,j} \otimes h_{1,2}^{i,j}$. If $m \in M$ then $\rho(m) = \sum m_0^i \otimes m_1^i$.

It is known (see e.g. [4] p.15) that for any Hopf algebra H and Hopf module M , $M \approx M' \otimes H$, where $M' = \{m \in M | \rho(m) = m \otimes 1\}$ with the isomorphism $m \mapsto \sum m_0^i \cdot S(m_{1,1}^{i,j}) \otimes m_{1,2}^{i,j}$, where this is actually a double sum on both i and j . It should also be noted that $M' \otimes H$ is a trivial Hopf module, i.e. one for which $(m \otimes h)l = m \otimes hl$.

If we now also tensor with k over H we will get

$$M \otimes_H k \approx (M' \otimes H) \otimes_H k.$$

However, since $M' \otimes H$ is a trivial (in the sense defined above) Hopf module we get

$$M \otimes_H k \approx (M' \otimes H) \otimes_H k \approx M' \otimes (H \otimes_H k) \approx M'.$$

The isomorphism is

$$m \otimes 1 \mapsto \sum m_0^i \cdot S(m_{1,1}^{i,j}) \otimes m_{1,2}^{i,j} \otimes 1 \mapsto \sum m_0^i \cdot S(m_{1,1}^{i,j}) \epsilon(m_{1,2}^{i,j}) = \sum m_0^i \cdot S(m_1^i).$$

If we take $M = W \otimes H$ with W any Hopf module, H acting with the diagonal action and

$$\rho(w \otimes h) = w \otimes \Delta(h)$$

then $M' = W \otimes k \approx W$. In this case, if $m = w \otimes h$ then $\rho(w \otimes h) = w \otimes \Delta(h)$ so $m_0^i = w \otimes h_1^i$ and $m_1^i = h_2^i$. Therefore the explicit form of the isomorphism is

$$w \otimes h \otimes 1 \mapsto \sum (w \otimes h_1^i) \Delta(S(h_2^i)).$$

However, we know that the image is in M' , so we can apply $1 \otimes \epsilon$ to the image and not change it. Also if $h \in H$ then from the definition of a Hopf algebra $(1 \otimes \epsilon)(\Delta(h)) = h \otimes 1$

Therefore the image is

$$(1 \otimes \epsilon)[\sum (w \otimes h_1^i) \Delta(S(h_2^i))] = \sum (w \otimes \epsilon(h_1^i))[(1 \otimes \epsilon)(\Delta(S(h_2^i)))] = \sum (w \otimes 1)(\epsilon(h_1^i) S(h_2^i) \otimes 1) = (w \otimes 1)(S(h) \otimes 1)$$

so the image in W is

$$w \otimes h \otimes 1 \mapsto w S(h).$$

In our case we are interested in the module $\otimes^n H$, so we can take $W = \otimes^{n-1} H$ and the isomorphism will be

$$h_1 \otimes h_2 \otimes \cdots \otimes h_n \otimes 1 \mapsto (h_1 \otimes h_2 \otimes \cdots \otimes h_{n-1}) \Delta_{n-1}(S(h_n)).$$

5 Computations

We can now prove theorem 1.1, i.e. show that $H_2(L/I^n, k)$ is not finitely generated by exhibiting an infinite number of elements of the multiplier, whose images in $\otimes^{n-1} U(L/I)$ are linearly independent. We shall deal with several cases. In each of them we

shall construct elements of $H_2(L/I^n, k)$ that have one parameter l , where $l \in U(L/I)$. In other words we shall construct a k -linear map $f : U(L/I) \rightarrow H_2(L/I^n, k) \rightarrow \otimes^{n-1} U(L/I)$. It is obviously enough to show that $\ker f = k \cdot 1$ (since $U(L/I)$ is not finite dimensional). In other cases we shall show that $\text{Im} f$ is not finite dimensional by proving that it has elements of unbounded degree.

Recall the elements $\alpha \in I/I'$ and $x \in X$ such that $a = \phi_x(\alpha)$ was non-zero, and consider all elements of the form $[\alpha \cdot l, \alpha, \dots, \alpha] \otimes 1$, where l is any element of $\delta U(L/I)$. Obviously this element is in I^n . Its image, using the mapping $\phi_{x,x,\dots,x}$ will be $[al, a, \dots, a] \otimes 1$. In other words $f(l) = [al, a, \dots, a] \otimes 1$. Note that if $l \in k \cdot 1$ then $f(l) = 0$ since in that case $[a \cdot l, a] = 0$. An easy induction shows that

$$[a, b, b, \dots, b] \otimes 1 = \sum (-1)^i \binom{n-1}{i} \otimes^i b \otimes a \otimes^{n-1-i} b \otimes 1$$

where $\otimes^i b$ means $b \otimes b \otimes \dots \otimes b$ (i times). The referee points out that this formula is known as the Cartan-Weyl formula. Therefore under the Hopf module isomorphism

$$\begin{aligned} f(l) &= \sum (-1)^i \binom{n-1}{i} (\otimes^i a \otimes al \otimes^{n-2-i} a) \Delta_{n-1}(S(a)) \\ &\quad + (-1)^{n-1} (\otimes^{n-1} a) \Delta_{n-1}(S(al)). \end{aligned}$$

But $S(al) = S(l)S(a)$ so $\Delta_{n-1}(S(al)) = \Delta_{n-1}(S(l))\Delta_{n-1}(S(a))$ and hence

$$\begin{aligned} f(l) &= [\sum (-1)^i \binom{n-1}{i} (\otimes^i a \otimes al \otimes^{n-2-i} a) \\ &\quad + (-1)^{n-1} (\otimes^{n-1} a) \Delta_{n-1}(S(l))] \Delta_{n-1}(S(a)). \end{aligned}$$

This can be rewritten as

$$\begin{aligned} f(l) &= (a \otimes a \otimes \dots \otimes a) [\sum (-1)^i \binom{n-1}{i} (\otimes^i 1 \otimes l \otimes^{n-2-i} 1) + \\ &\quad (-1)^{n-1} \Delta_{n-1}(S(l))] \Delta_{n-1}(S(a)). \end{aligned}$$

Since $U(L/I)$ is without zero divisors and we are only interested in $\ker f$ or the dimension of $\text{Im} f$, we can consider instead the function

$$f(l) = \sum (-1)^i \binom{n-1}{i} (\otimes^i 1 \otimes l \otimes^{n-2-i} 1) + (-1)^{n-1} \Delta_{n-1}(S(l)).$$

In order to compute $\ker f$, we can apply ϵ to all but the j -th coordinate of each monomial. This operator, applied to $\otimes^i 1 \otimes l \otimes^{n-2-i} 1$, yields $l\delta_{ij}$ (since $\epsilon(l) = 0$), while applied to $\Delta_{n-1}(S(l))$ yields (because ϵ is a counit) $S(l)$. Therefore for each $0 \leq j < n$ the result is

$$(-1)^j \binom{n-1}{j} l + (-1)^{n-1} S(l) = 0.$$

Therefore $S(l) = (-1)^{n+j} \binom{n-1}{j} l$.

If $n > 2$ we get $S(l) = (-1)^n l$ and $S(l) = (-1)^{n+1}(n-1)l$.

Therefore $(-1)^n l = (-1)^{n+1}(n-1)l$ i.e.

$$nl = 0.$$

As was mentioned above, there are several cases.

Case I If $\text{char}(k)$ does not divide n and $n > 2$ then for any $l \in \delta U(L/I)$ we have $f(l) \neq 0$ i.e. $\ker f = k \cdot 1$.

Case II If $\text{char}(k) \neq 2$. We wish to show that $\text{Im} f$ is not finite dimensional. Denoting by $f_1(l)$ the application of ϵ to all but the first coordinate, we get $f_1(l) = l + (-1)^{n-1} S(l)$. This is true also when $n = 2$. Since f_1 is simply f composed with another function, obviously $\dim(\text{Im} f_1) \leq \dim(\text{Im} f)$. Therefore it is enough to consider f_1 . However, if x is any non-zero Lie element in $U(L/I)$ then $S(x^i) = (-1)^i x^i$. So $f_1(x^i) = x^i + (-1)^{i+n-1} x^i$. Since $\text{char}(k) \neq 2$ then for all i of the correct parity we will have $f_1(x^i) = 2x^i \neq 0$, but $\deg x^i = i$ will be unbounded, so we are finished.

Case III The only case left is $\text{char}(k) = 2$ and n even. In this case we still have $f_1(l) = l - S(l)$. Suppose L/I is not commutative, therefore there exist $x, y \in L$ such that $[x, y] \notin I$, i.e. $[x, y] \neq 0$ in $U(L/I)$. Consider $l_i = xy^i$. Obviously $S(l_i) = y^i x$, so $f_1(l_i) = xy^i - y^i x = [x, y^i]$. However the mapping $u \mapsto [x, u]$ is a derivation of $U(L/I)$, and therefore

$$[x, y^i] = \sum_{j=0}^{i-1} y^j [x, y] y^{i-j-1}.$$

Note that $[x, y]y = y[x, y] + [[x, y], y]$, and hence $y^j [x, y] y^{i-j-1} \equiv y^{i-1} [x, y] \pmod{U_{i-1}(L/I)}$. Thus $[x, y^i] \equiv i y^{i-1} [x, y] \pmod{U_{i-1}(L/I)}$, and if i is odd then $\deg f_1(l_i) = i$. Thus the degree of the elements of the image is unbounded, so the image is infinite dimensional.

Case IV There remains the case where L/I is commutative. Thus if L has basis X , then $L' \subset I$ so $I/L' \subset L/L'$ is a subspace, and we

can perform a linear change of basis of L , so that $I = \langle L', X_1 \rangle$, where X_1 is a proper subset of X . Consider the Lie algebra over $\mathbb{Z}$, $L_1 = \langle Y \rangle$, $I_1 = \langle L'_1, Y_1 \rangle$, where Y and Y_1 are disjoint copies of X and X_1 . We now use the universal coefficient theorem (see e.g. [3] p.176) which in our case states that if k is any $\mathbb{Z}$ module then

$$\begin{aligned} 0 \rightarrow H_2(L_1/I_1^n, \mathbb{Z}) \otimes_{\mathbb{Z}} k &\rightarrow H_2(L_1/I_1^n \otimes_{\mathbb{Z}} k, k) \\ &\rightarrow \text{Tor}_1^{\mathbb{Z}}(H_1(L_1/I_1^n, \mathbb{Z}), k) \rightarrow 0 \end{aligned}$$

is exact. Since $H_1(L_1/I_1^n, \mathbb{Z}) = (L_1/I_1^n)_{ab} = L_1/L'_1$ is a free $\mathbb{Z}$ module then $\text{Tor}_1^{\mathbb{Z}}(H_1(L_1/I_1^n, \mathbb{Z}), k) = 0$.

Take $k = \mathbb{Q}$. We have $H_2(L_1/I_1^n, \mathbb{Z}) \otimes_{\mathbb{Z}} \mathbb{Q} \approx H_2(L_1/I_1^n \otimes_{\mathbb{Z}} \mathbb{Q}, \mathbb{Q})$. However $L_1/I_1^n \otimes_{\mathbb{Z}} \mathbb{Q}$ is simply L_2/I_2^n where $L_2 = \langle Y \rangle$ and $I_2 = \langle L'_2, Y_1 \rangle$ taken over $\mathbb{Q}$. Since $\mathbb{Q}$ has characteristic 0, we know that $H_2(L_2/I_2^n \otimes_{\mathbb{Z}} \mathbb{Q}, \mathbb{Q})$ is infinite dimensional. Therefore $H_2(L_1/I_1^n, \mathbb{Z})$ must also have infinite torsion-free rank as a $\mathbb{Z}$ -module. Apply now the universal coefficient theorem with k any field of characteristic 2. Again $H_2(L_1/I_1^n, \mathbb{Z}) \otimes_{\mathbb{Z}} k \approx H_2(L_1/I_1^n \otimes_{\mathbb{Z}} k, k)$. Once again $L_1/I_1^n \otimes_{\mathbb{Z}} k$ is exactly L/I^n of the original Lie algebra. However, since $H_2(L_1/I_1^n, \mathbb{Z})$ has infinite rank then $H_2(L_1/I_1^n, \mathbb{Z}) \otimes_{\mathbb{Z}} k$ is not finitely generated, thus we have proved theorem 1.1.

Note that in the case $L = \langle x, y \rangle$, $I = L'$ and k is of characteristic 2, even though $H_2(L/I', k)$ is not finitely generated, the image in $U(L/I)$, under any of the projections, will be 0.

References

- [1] G. Baumslag, R. Strebel and M Thomson, On the Multiplier of $F/\gamma_c R$, *J. Pure Appl. Alg.* **16** (1980) 121-132.
- [2] G. Bergman and W. Dicks, On Universal Derivations, *J. Alg.* **36** (1975) 193-211.
- [3] P.J. Hilton and U. Stammach, A course in Homological Algebra, *Graduate Texts in Mathematics* **4**, Springer-Verlag, 1971.
- [4] S. Montgomery, Hopf Algebras and Their Actions on Rings, *Regional Conference Series in Mathematics* **82**, American Mathematical Society, 1992.
- [5] J. Lewin, On Some Infinitely Presented Associative Algebras, *J. Austr. Math. Soc.* **16** (1973) 290-293.

- [6] C. Reutenauer, Free Lie Algebras, *London Math. Soc. Monographs New Series* **7**, Oxford Univ. Press, 1993.
- [7] C. Weibel, An Introduction to Homological Algebra, *Cambridge Studies in Advanced Math.* **38**, Cambridge Univ. Press, 1994.